Fluid Phase Equilibria, 47 (1989) 311–312
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands

Contents Volume 47 (1989)

Volume 47, No. 1

Prediction of ternary phase equilibria by the PRSV2 equation of state with the NRTL mixing rule
T. Ohta (Kanazawa, Japan) 1– 15
New equation of state based on the significant structure model
K. Suzuki, H. Sue (Chiba, Japan), R.L. Smith, H. Inomata, K. Arai and S. Saito (Sendai, Japan) 17– 38
On the correspondence between the density functions and the partial molar volume functions in binary mixtures
Y.P. Puhovski and Y.M. Kessler (Ivanovo, U.S.S.R.) 39– 46
Application of the group contribution concept to the Kihara potential for the calculation of second virial coefficients
S.W. Campbell (Tampa, FL, U.S.A.) 47– 65
Determination of the coexistence curve of sulfur hexafluoride from isochoric intercepts
S.N. Biswas and C.A. ten Seldam (Amsterdam, The Netherlands) 67– 75
A density correction for the Peng–Robinson equation of state
P.M. Mathias, T. Naheiri and E.M. Oh (Allentown, PA, U.S.A.) 77– 87
Excess enthalpies of binary mixtures of benzene with n-propylbenzene, n-butylbenzene, n-hexylbenzene at 298.15, 308.15 and 318.15 K
D.V.S. Jain and N.S. Dhar (Chandigarh, India) 89– 94
Excess molar volumes of ethylacetate + hydrocarbons at 303.15 K: an interpretation in terms of the Prigogine–Flory–Patterson theory
A.M. Awwad, K.A. Jabra and A.H. Al-Dujaili (Baghdad, Iraq) 95–102

Short Communication
Mixing rules for hydrogen-containing systems
W. Wang and C. Zhong (Beijing, China) 103–114

Volume 47, Nos. 2–3

Excess functions and equations of state
A. Peneloux, W. Abdoul and E. Rauzy (Marseille, France) 115–132
Prediction of gas solubility and vapor–liquid equilibria by group contribution
R. Gani, N. Tzouvaras, P. Rasmussen and A.A. Fredenslund (Lyngby, Denmark) 133–152
Development and application of Simha equation of state to calculation of the density of certain alkanes and their mixtures
H. Zhou, B. Lagourette, J. Alliez, P. Xans and F. Montel (Pau, France) 153–169
A new pseudocubic perturbed hard-sphere equation of state
M. Kato, M. Yamaguchi and T. Kiuchi (Fukushima, Japan) 171–187

A modified Peng–Robinson equation of state
G.A. Melhem (Cambridge, MA, U.S.A.), R. Saini and B.M. Goodwin (Boston, MA, U.S.A.) 189–237

On the importance of the first order blip expansion term in WCA-type perturbation theories
J. Fischer, A. Lotfi (Bochum, F.R.G.), K. Lucas (Duisburg, F.R.G.) and K.P. Shukla (Tulsa, OK, U.S.A.) 239–248

Thermodynamics of liquid mixtures of xenon and methyl fluoride
I.M.A. Fonseca and L.Q. Lobo (Coimbra, Portugal) 249–263

Gas–liquid pressure–temperature–composition critical loci for propanenitrile with C_5–C_{11} n-alkanes
A. Trejo and M.C. Zepeda (Mexico, D.F., Mexico) 265–271

Apparatus and procedure to measure binary total pressure at high temperature fluorene–phenanthrene vapor–liquid equilibria and data reduction by the four-suffix Margules model to obtain infinite-dilution activity coefficients
V. Roper and R. Kobayashi (Houston, TX, U.S.A.) 273–293

Final hydrate stability conditions of a methane and propane mixture in the presence of pure water and aqueous solutions of methanol and ethylene glycol
K.Y. Song and R. Kobayashi (Houston, TX, U.S.A.) 295–308

Author Index 309

Contents of Volume 47 (1989) 311–312